

The case indicates that although bilateral primary renal lymphoma is a rare clinical condition, but sometimes it is possible.

### Conclusion

If a lymphomatous malignancy is found in core biopsy of a renal mass, nephrectomy can be avoided and the patient can be treated with systemic chemotherapy. At present the CHOP regimen is considered to be a standard treatment for renal lymphoma, but only 41% patients with advanced disease are alive and disease free at a median follow up of 3 years.

### References

1. Ajeet Gajra Md et all "Lymphoma;B cell". eMedicine Journal [serial online].2005. Available at: <http://www.e-medicine.com/med/topic1358.htm>
2. Pddubnaya,Y.Balakireva,D. Osmanov .Prognostic factors in primary extranodal non-Hodgkin's lymphoma patients.5.Extranodal lymphomas. Ann. Onc., Jun 2005;16:v159-v167
3. Ali Shirkhoda, MD, FACR. "Kidney,lymphoma". eMedicine Journal [serial online]. 2005. Available at: <http://www.e-medicine.com/radio/topic373.htm>
4. Alan S.L. Yu,Barry M.Brenner: Tubulo - interstitial disease of kidney.In Dennis L kasper MD,Eugene Braunwald MD,Anthony S Fauci MD,Stephen L Hauser MD, Dan L. Longo MD,J Larry Jameson MD,Phd ed. Harrison's Principles of Internal. Medicine.16th edition. 2005. McGraw-Hill,USA. Pp 1704-1705
5. Hassan Pervez, Muhammad Shaikh, Anil Pottis, , Sayeed A.Mehdi. Diagnosis in oncology; Uncommon presentation of Non-Hodgkin's lymphoma. J Clin Oncol,2003 Feb ; 21(3): 567- 569
6. Nelson G. Ordonez and Juan Rosai:Urinary Tract: Kidney, renal pelvis and ureter.In Juan Rosai MD ed. Rosai and Ackerman's Surgical Pathology Volume I. thedition. 2004, Mosby, Elsevier,India. Pp-1272
7. James O.Armitage, Dan L.Longo : Malignancies of lymphoid cells. In Dennis L kasper MD, Eugene Braunwald MD, Anthony S Fauci MD, Stephen L Hauser MD, Dan L. Longo MD,J Larry Jameson MD,Phd ed. Harrison's Principles of Internal Medicine.16th edition. 2005. McGraw-Hill, USA.Pp 649-650

□